

From vacuum field equations on principal bundles to Einstein's equations with fluids

M. MONTESINOS-VELÁSQUEZ* AND T. MATOS†

Departamento de Física

Centro de Investigación y de Estudios Avanzados del IPN

Apartado Postal 14-740, 07000, México, D.F., México

ABSTRACT. In the present work we show that the Einstein equations on M without cosmological constant and with perfect fluid as source, can be obtained from the field equations for vacuum with cosmological constant on the principal fibre bundle $P\left(\frac{1}{I}M, U(1)\right)$, M being the space-time and I the radius of the internal space $U(1)$.

RESUMEN. Mostramos que las ecuaciones de Einstein sobre M sin constante cosmológica y con fluido perfecto como fuente, pueden obtenerse a partir de las ecuaciones de campo para vacío con constante cosmológica sobre el haz fibrado principal $P\left(\frac{1}{I}M, U(1)\right)$, donde M es el espacio-tiempo e I el radio del espacio interno $U(1)$.

PACS: 11.10.Kk ; 98.80.Dr

*Electronic address : merced@fis.cinvestav.mx

†Electronic address : tmatos@fis.cinvestav.mx

1. INTRODUCTION

In a recent work [1] it has been shown that vacuum solutions in scalar-tensor theories are equivalent to solutions of general relativity with imperfect fluid as source. The above models have the defect that the scalar fields do not arise from a natural framework of unification, but they are put by hand as in the inflationary models [2] and are therefore artificial fields in the theory. On the other hand, we know that the geometric formalism of principal fibre bundles [3,4] is a natural scheme to unify the general relativity theory with gauge field theories (Abelian and Non-Abelian). If the principal fibre bundle $P(\tilde{M}, U(1))$ is endowed with a metric “dimensionally reducible” to $\tilde{M}$ by means of the reduction theorem [5], *i.e.*, if the metric can be built out from quantities defined only on M , then the scalar fields arise in a natural way. Therefore, it is important to study the above model in the context of [1] for the particular principal fibre bundle $P(\frac{1}{7}M, U(1))$, M being the space-time and I the scalar field. This paper is organized as follows: in the next section we review the geometric formalism of principal fibre bundles while in Sect. 3 we deduce the Einstein equations without cosmological constant and perfect fluid as source from the field equations on $P(\frac{1}{7}M, U(1))$ for vacuum and cosmological constant. We give an example in Sect. 4 when $\tilde{M}$ is conformally *FRW*. Finally we summarize the results in Sect. 5.

2. THE GEOMETRY

The actual version of the Kaluza-Klein theories is based on the mathematical structure of principal fibre bundles [5,6]. In this scheme, the unification of the general relativity theory with the gauge theories is a natural fact. Moreover, the reduction theorem provides a metric on (right) principal fibre bundles $P(\tilde{M}, G)$ which is right-invariant under the action of the structure group G on the whole space P . In the trivialization of the bundle this metric

reads [5,6]

$$\hat{g} = \tilde{g}_{\alpha\beta} dx^\alpha \otimes dx^\beta + \xi_{mn} (\omega^m + A_\alpha^m dx^\alpha) \otimes (\omega^n + A_\beta^n dx^\beta), \quad (1)$$

where the metric of the base space $\tilde{M}$ (generally identified with the space-time of general relativity) is $\tilde{g}_{\alpha\beta} dx^\alpha \otimes dx^\beta$ while the metric on the fibre ($x^\alpha = \text{const.}$) is $\xi_{mn} \omega^m \otimes \omega^n$ and $\{\omega^m\}$ is a basis of right-invariant 1-forms on G . The quantities $\tilde{g}_{\alpha\beta}$, ξ_{mn} and A_α^n depend only on the coordinates on $\tilde{M}$ and the A_α^n correspond to Yang-Mills potentials in the gauge theory while the ξ_{mn} are the scalar fields.

In particular, the principal fibre bundle $P(\tilde{M}, U(1))$ has the metric

$$\hat{g} = \tilde{g}_{\alpha\beta} dx^\alpha \otimes dx^\beta + I^2 (d\psi + A_\alpha dx^\alpha) \otimes (d\psi + A_\beta dx^\beta), \quad (2)$$

where the scalar field I correspond to the radius of the internal space $U(1)$ and ψ is the coordinate on $U(1)$ too. However, the magnitude of the internal radius I depends on the particular cases; cosmological or astrophysical models (for details on units and magnitude on the scalar field I see Refs. [6,7]). For vanishing electromagnetic potential, $A_\alpha = 0$, we obtain the unification of $\tilde{g}_{\alpha\beta}$ with the scalar field I

$$\hat{g} = \tilde{g}_{\alpha\beta} dx^\alpha \otimes dx^\beta + I^2 d\psi^2. \quad (3)$$

By using Eq. (3) we compute the Ricci tensor

$$\hat{R}_{\alpha\beta} = \tilde{R}_{\alpha\beta} - I^{-1} I_{;\alpha\beta}, \quad (4)$$

$$\hat{R}_{\alpha 4} = 0, \quad (5)$$

$$\hat{R}_{44} = -I \square I, \quad (6)$$

where greek indices run on 0,1,2,3 and the label “4” corresponds to the fifth dimension.

Usually the base space $\tilde{M}$ of $P(\tilde{M}, G)$ is identified as the space-time, in this paper we adopt the version where the base space $\tilde{M}$ of $P(\tilde{M}, U(1))$ is conformally the space-time M

of general relativity, *i.e.*, $\tilde{M} = \frac{1}{7}M$. That is to say, we start with the metric (compare Ref. [8])

$$\hat{g} = \frac{1}{I} g_{\alpha\beta} dx^\alpha \otimes dx^\beta + I^2 d\psi^2, \quad (7)$$

where $g_{\alpha\beta} dx^\alpha \otimes dx^\beta$ is the space-time metric. Then by using Eq. (7) we obtain the Ricci tensor

$$\hat{R}_{\alpha\beta} = R_{\alpha\beta} + \frac{1}{2} \left(I^{-1} \square I - I_{;\lambda} I^{;\lambda} \right) g_{\alpha\beta} - \frac{3}{2} I^{-2} I_{;\alpha} I_{;\beta}, \quad (8)$$

$$\hat{R}_{\alpha 4} = 0, \quad (9)$$

$$\hat{R}_{44} = -I^2 \square I + I_{;\lambda} I^{;\lambda}. \quad (10)$$

In the next we use the signature $(-, +, +, +)$ for the space-time metric on M .

3. PERFECT FLUID STRUCTURE

The field equations on $P(\frac{1}{7}M, U(1))$ in vacuum with cosmological constant Λ are given by $\hat{R}_{AB} - \frac{\hat{R}}{2} \hat{g}_{AB} = \Lambda \hat{g}_{AB}$ or in equivalent form

$$\hat{R}_{AB} = -\frac{2}{3} \Lambda \hat{g}_{AB}, \quad (11)$$

where A, B run on greek indices α and 4.

By using Eqs. (8)-(10) and (11) we obtain

$$R_{\alpha\beta} = I^{-2} \left(\frac{1}{2} I_{;\lambda} I^{;\lambda} g_{\alpha\beta} + \frac{3}{2} I_{;\alpha} I_{;\beta} \right) - I^{-1} \left(\frac{1}{2} \square I + \frac{2}{3} \Lambda \right) g_{\alpha\beta}, \quad (12)$$

$$\square I = \frac{1}{I} I_{;\lambda} I^{;\lambda} + \frac{2}{3} \Lambda. \quad (13)$$

By substituting the field equation for I [Eq. (13)] into the Ricci tensor [Eq. (12)] we obtain the next equivalent equations system

$$R_{\alpha\beta} = \frac{3}{2} I^{-2} I_{;\alpha} I_{;\beta} - I^{-1} \Lambda g_{\alpha\beta}, \quad (14)$$

$$\square I = \frac{1}{I} I_{;\lambda} I^{;\lambda} + \frac{2}{3} \Lambda. \quad (15)$$

On the other hand, by using the Einstein equations without cosmological constant,

$$R_{\alpha\beta} = T_{\alpha\beta} - \frac{T}{2}g_{\alpha\beta} \quad (16)$$

and Eq. (14), we can define the energy-momentum tensor associated with the scalar field I

$$T_{\alpha\beta} = \frac{3}{2}I^{-2}I_{;\alpha}I_{;\beta} + \left(-\frac{3}{4}I^{-2}I_{;\lambda}I^{;\lambda} + I^{-1}\Lambda\right)g_{\alpha\beta}. \quad (17)$$

This energy-momentum tensor is covariantly conserved, $T^{\alpha\beta}_{;\beta} = 0$ as follows from the field equation for I . Finally, by comparing the above energy-momentum tensor associated with the scalar field I with that of an imperfect fluid

$$T_{\alpha\beta} = \rho U_\alpha U_\beta + 2q_{(\alpha}U_{\beta)} + p h_{\alpha\beta} + \pi_{\alpha\beta}, \quad (18)$$

where ρ is the energy density of fluid, U_α the velocity, q_α the heat flux vector, p the pressure, $\pi_{\alpha\beta}$ the anisotropic stress tensor and

$$h_{\alpha\beta} = g_{\alpha\beta} + U_\alpha U_\beta, \quad (19)$$

is the projection orthogonal to the velocity, we conclude [1]

$$q_\alpha = 0, \quad (20)$$

$$\pi_{\alpha\beta} = 0, \quad (21)$$

$$\rho = -\frac{3}{4}I^{-2}I_{;\lambda}I^{;\lambda} - \frac{\Lambda}{I}, \quad (22)$$

$$p = -\frac{3}{4}I^{-2}I_{;\lambda}I^{;\lambda} + \frac{\Lambda}{I}, \quad (23)$$

where the velocity has been choosen in the form [1]

$$U_\alpha \equiv \frac{I_{;\alpha}}{\sqrt{-I_{;\lambda}I^{;\lambda}}} \quad (24)$$

That is to say, Eqs. (20)-(23) implies that Eq. (17) has the structure corresponding to a perfect fluid. Moreover, if $\Lambda = 0$ then Eqs. (17) and (20)-(23) correspond to the so called ‘‘Zeldovich ultrastiff matter’’ fluid, $p = \rho$ (see Ref. [1]).

4. EXAMPLE: THE P.F.B. $P(\frac{1}{I}FRW, U(1))$

We start from the metric

$$\hat{g} = \frac{1}{I(t)} \left[-dt^2 + R^2(t) \left(\frac{dr^2}{1 - k r^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right) \right] + I^2(t) d\psi^2, \quad (25)$$

where $I = I(t)$ on account of the isotropy and homogeneity of the FRW metric. In this case the Eqs. (14)-(15) read

$$-3 \left(\frac{\ddot{R}}{R} \right) = \frac{3}{2} \left(\frac{\dot{I}}{I} \right)^2 + \left(\frac{1}{I} \right) \Lambda, \quad (26)$$

$$2 \left(\frac{k}{R^2} \right) + 2 \left(\frac{\dot{R}}{R} \right)^2 + \left(\frac{\ddot{R}}{R} \right) = - \left(\frac{1}{I} \right) \Lambda, \quad (27)$$

$$3 \left(\frac{\dot{R}}{R} \right) \left(\frac{\dot{I}}{I} \right) - \left(\frac{\dot{I}}{I} \right)^2 + \left(\frac{\ddot{I}}{I} \right) = - \frac{2}{3} \left(\frac{1}{I} \right) \Lambda, \quad (28)$$

where dot means derivation with respect to the cosmological time t . These equations are equivalent to the Einstein equations for FRW with perfect fluid as source, provided that

$$\rho = \frac{3}{4} \left(\frac{\dot{I}}{I} \right)^2 - \left(\frac{1}{I} \right) \Lambda, \quad (29)$$

$$p = \frac{3}{4} \left(\frac{\dot{I}}{I} \right)^2 + \left(\frac{1}{I} \right) \Lambda. \quad (30)$$

By the way, the field equation for I [Eq. (28)] is the covariant conservation of $T_{\alpha\beta}$, $T^{\alpha\beta}_{;\beta} = 0$

$$\dot{\rho} + 3 \left(\frac{\dot{R}}{R} \right) (\rho + p) = 0. \quad (31)$$

5. CONCLUSION

We have shown that the field equations with cosmological constant Λ on the principal fibre bundle $P(\frac{1}{I}M, U(1))$ are equivalent to the Einstein equations without cosmological

constant on M and with perfect fluid as source. In order to show it we start from the field equations on $P(\frac{1}{I}M, U(1))$, $\hat{R}_{AB} = -\frac{2}{3}\Lambda \hat{g}_{AB}$ and separate them in their 4-dimensional and five dimensional parts. We have found that from the 4-dimensional part of these equations it is possible to define an effective energy-momentum tensor $T_{\alpha\beta}$ and that it is covariantly conserved, being $T^{\alpha\beta}_{;\beta} = 0$ equivalent to the field equation for I . Finally, we applied the above result to the particular bundle $P(\frac{1}{I}FRW, U(1))$.

ACKNOWLEDGMENTS

The authors are grateful to CONACyT for financial support.

REFERENCES

1. L.O. Pimentel, *Class. Quantum Grav.* **6** (1989) L263.
2. E.W. Kolb and M.S. Turner, *The Early Universe*, Addison-Wesley, Redwood City, CA (1990).
3. A. Trautman, *Rep. Math. Phys.* **1** (1970) 29.
4. Y. Choquet-Bruhat, C. De Witt-Morette and the M. Dillard-Bleick, *Analysis, Manifolds and Physics*, Part I, Revised Edition, North-Holland (1991).
5. C. Choquet-Bruhat and C. De Witt-Morette, *Analysis, Manifolds and Physics*, Part II, North-Holland (1989), Sect. V.13.
6. T. Matos and J.A. Nieto, *Rev. Mex. Fís.* **39** No. Sup. 1 (1993) 81.
7. D. Bailin and A. Love, *Rep. Prog. Phys.* **50** (1987) 1087.
8. P.S. Wesson and J. Ponce de Le'ón, *J. Math. Phys.* **33** (11) (1992) 3883.